



ELSEVIER



C. R. Acad. Sci. Paris, Ser. I ●●● (●●●●) ●●●-●●●



COMPTES RENDUS

MATHÉMATIQUE

http://france.elsevier.com/direct/CRASS1/

## Group Theory

# On the number of conjugates defining the solvable radical of a finite group

Nikolai Gordeev<sup>a</sup>, Fritz Grunewald<sup>b</sup>, Boris Kunyavskii<sup>c</sup>, Eugene Plotkin<sup>c</sup>

<sup>a</sup> Department of Mathematics, Herzen State Pedagogical University, 48 Moika Embankment, 191186 St. Petersburg, Russia

<sup>b</sup> Mathematisches Institut der Heinrich-Heine-Universität Düsseldorf, Universitätsstr. 1, 40225 Düsseldorf, Germany

<sup>c</sup> Department of Mathematics, Bar-Ilan University, Ramat Gan 52900, Israel

Received 12 July 2006; accepted 23 August 2006

Presented by Jean-Pierre Serre

## Abstract

We are looking for the smallest integer  $k > 1$  providing the following characterization of the solvable radical  $R(G)$  of any finite group  $G$ :  $R(G)$  consists of the elements  $g$  such that for any  $k$  elements  $a_1, a_2, \dots, a_k \in G$  the subgroup generated by the elements  $g, a_i g a_i^{-1}, i = 1, \dots, k$ , is solvable. Our method is based on considering a similar problem for commutators: find the smallest integer  $\ell > 1$  with the property that  $R(G)$  consists of the elements  $g$  such that for any  $\ell$  elements  $b_1, b_2, \dots, b_\ell \in G$  the subgroup generated by the commutators  $[g, b_i], i = 1, \dots, \ell$ , is solvable. **To cite this article:** N. Gordeev et al., C. R. Acad. Sci. Paris, Ser. I ●●● (●●●●).

© 2006 Published by Elsevier SAS on behalf of Académie des sciences.

## Résumé

**Sur le nombre de conjugués définissant le radical résoluble d'un groupe fini.** Nous cherchons le plus petit entier  $k > 1$  caractérisant le radical résoluble  $R(G)$  d'un groupe fini  $G$  comme suit :  $R(G)$  est l'ensemble des éléments  $g$  tels que pour toute partie à  $k$  éléments  $\{a_1, a_2, \dots, a_k\} \subset G$  le sous-groupe engendré par les éléments  $g, a_i g a_i^{-1}, i = 1, \dots, k$ , est résoluble. Notre méthode s'appuie sur la considération d'un problème similaire pour les commutateurs. Nous cherchons le plus petit entier  $\ell > 1$  ayant la propriété suivante :  $R(G)$  est l'ensemble des éléments  $g$  tels que pour toute partie à  $\ell$  éléments  $\{b_1, b_2, \dots, b_\ell\} \subset G$  le sous-groupe engendré par les commutateurs  $[g, b_i], i = 1, \dots, \ell$ , est résoluble. **Pour citer cet article :** N. Gordeev et al., C. R. Acad. Sci. Paris, Ser. I ●●● (●●●●).

© 2006 Published by Elsevier SAS on behalf of Académie des sciences.

## Version française abrégée

Notre but est une nouvelle caractérisation du radical résoluble  $R(G)$  d'un groupe fini  $G$ .

**Théorème 0.1.** *Le radical résoluble d'un groupe fini  $G$  est l'ensemble des éléments  $g$  satisfaisant la propriété suivante : pour toute partie à 7 éléments  $\{a_1, a_2, \dots, a_7\} \subset G$ , le sous-groupe engendré par les éléments  $g, a_i g a_i^{-1}, i = 1, \dots, 7$ , est résoluble.*

*E-mail addresses:* nickgordeev@mail.ru (N. Gordeev), grunewald@math.uni-duesseldorf.de (F. Grunewald), kunyav@macs.biu.ac.il (B. Kunyavskii), plotkin@macs.biu.ac.il (E. Plotkin).

1631-073X/\$ – see front matter © 2006 Published by Elsevier SAS on behalf of Académie des sciences.

doi:10.1016/j.crma.2006.08.005

1 La démonstration utilise la classification des groupes simples finis.

2 Cet énoncé implique la caractérisation suivante des groupes résolubles finis :

3  
4 **Théorème 0.2.** *Un groupe fini  $G$  est résoluble si et seulement si dans chaque classe de conjugaison de  $G$  toute partie*  
5 *à 8 éléments engendre un sous-groupe résoluble.*

6  
7 Nous espérons d'améliorer ces caractérisations.

8  
9 **Conjecture 0.3.** *Le radical résoluble d'un groupe fini  $G$  est l'ensemble des éléments  $g$  satisfaisant la propriété*  
10 *suivante : pour toute partie à trois éléments  $\{a, b, c\} \subset G$ , le sous-groupe engendré par les éléments  $g, aga^{-1}, bgb^{-1},$*   
11  *$cgc^{-1}$  est résoluble.*

12  
13 Cette assertion implique :

14  
15 **Conjecture 0.4.** *Un groupe fini  $G$  est résoluble si et seulement si dans chaque classe de conjugaison de  $G$  toute*  
16 *partie à quatre éléments engendre un sous-groupe résoluble.*

17  
18 **Remarque 0.5.** Ces caractérisations sont les meilleures possibles : dans les groupes symétriques  $S_n$  ( $n \geq 5$ ) trois  
19 transpositions engendrent un sous-groupe résoluble.

20  
21 **Remarque 0.6.** Le point clé dans notre démonstration du théorème 0.1 est le théorème 0.8 ci-dessous. Pour démontrer  
22 la conjecture 0.3 (et donc la conjecture 0.4), il faut étendre l'énoncé du théorème 0.8 à tous les groupes *presque simples*  
23 (c'est-à-dire, aux groupes  $H$  tels que  $G \subseteq H \subseteq \text{Aut}(G)$  pour certain groupe simple  $G$ ).

24  
25 **Définition 0.7.** Soit  $k \geq 2$  un nombre entier. On dit que  $g \in G$  est un élément  $k$ -radiciel si pour toute partie à  $k$   
26 éléments  $\{a_1, \dots, a_k\} \subset G$ , le sous-groupe engendré par les commutateurs  $[a_1, g], \dots, [a_k, g]$  est résoluble.

27  
28 Nous démontrons :

29  
30 **Théorème 0.8.** *Soit  $G$  un groupe simple fini non abélien. Alors  $G$  ne contient pas d'élément 3-radiciel non trivial.*

31  
32 Ce théorème implique les Théorèmes 0.1 and 0.2.

33  
34 La preuve consiste à faire une analyse cas par cas (groupes alternés, groupes de type de Lie, groupes sporadiques) ;  
35 dans certains cas on obtient un énoncé plus précis.

36  
37 **Définition 0.9.** Notons  $\rho(g)$  le plus petit entier  $n$  ayant la propriété suivante :  $g \notin R(G)$  si et seulement s'il existe  
38  $x_1, \dots, x_n \in G$  tels que le sous-groupe engendré par les commutateurs  $[g, x_1], \dots, [g, x_n]$  n'est pas résoluble. On  
39 appelle le nombre  $\rho(G) := \max_{g \in G \setminus R(G)} \rho(g)$  le degré radiciel de  $G$ .

40  
41 **Théorème 0.10.** *Si  $G$  est un groupe simple fini non abélien, alors  $\rho(G) \leq 3$ . Si  $G$  est un groupe de type de Lie sur*  
42 *un corps  $K$  tel que  $\text{char } K \neq 2$  et  $K \neq \mathbb{F}_3$ , ou un groupe sporadique non isomorphe à  $Fi_{22}$  ou  $Fi_{23}$ , alors  $\rho(G) = 2$ .*

## 43 44 1. Main results

45  
46 Our goal is to prove a new characterization of the solvable radical  $R(G)$  of a finite group  $G$ .

47  
48 **Theorem 1.1.** *The solvable radical of any finite group  $G$  consists of the elements  $g$  satisfying the property: for any 7*  
49 *elements  $a_1, a_2, \dots, a_7 \in G$  the subgroup generated by the elements  $g, a_i ga_i^{-1}, i = 1, \dots, 7$ , is solvable.*

50  
51 The proof involves the classification of finite simple groups.

52 This theorem implies the following characterization of finite solvable groups:

**Theorem 1.2.** *A finite group  $G$  is solvable if and only if in each conjugacy class of  $G$  every 8 elements generate a solvable subgroup.*

We hope to strengthen these results.

**Conjecture 1.3.** *The solvable radical of a finite group  $G$  consists of the elements  $g$  satisfying the property: for any 3 elements  $a, b, c \in G$  the subgroup generated by conjugates  $g, aga^{-1}, bgb^{-1}, cgc^{-1}$  is solvable.*

This statement implies:

**Conjecture 1.4.** *A finite group  $G$  is solvable if and only if in each nontrivial conjugacy class of  $G$  every four elements generate a solvable subgroup.*

**Remark 1.5.** These characterizations are the best possible: in the symmetric groups  $S_n$  ( $n \geq 5$ ) any triple of transpositions generates a solvable subgroup.

**Remark 1.6.** The main step in our proof of Theorem 1.1 is Theorem 1.8. To prove Conjecture 1.3 (and hence Conjecture 1.4), one has to extend the statement of Theorem 1.8 to all *almost simple* groups, i.e. to the groups  $H$  such that  $G \subseteq H \subseteq \text{Aut}(G)$  for some simple group  $G$  (see Remark 2.1 below).

Throughout the note  $\langle a_1, \dots, a_k \rangle$  stands for the subgroup of  $G$  generated by  $a_1, \dots, a_k \in G$ . We define the commutator of  $x, y \in G$  by  $[x, y] = xyx^{-1}y^{-1}$ .

**Definition 1.7.** Let  $k \geq 2$  be an integer. We say that  $g \in G$  is a *k-radical element* if for any  $a_1, \dots, a_k \in G$  the subgroup  $H = \langle [a_1, g], \dots, [a_k, g] \rangle$  is solvable.

We prove

**Theorem 1.8.** *Let  $G$  be a finite non-Abelian simple group. Then  $G$  does not contain nontrivial 3-radical elements.*

This theorem implies Theorems 1.1 and 1.2.

The proof goes by case-by-case inspection (alternating groups, groups of Lie type, sporadic groups); in some cases the statement is more precise.

**Definition 1.9.** Denote by  $\rho(g)$  the smallest possible  $n$  with the following property:  $g \notin R(G)$  if and only if there exist  $x_1, \dots, x_n \in G$  such that the subgroup  $\langle [g, x_1], \dots, [g, x_n] \rangle$  is not solvable. We call the number  $\rho(G) := \max_{g \in G \setminus R(G)} \rho(g)$  the radical degree of  $G$ .

**Theorem 1.10.** *If  $G$  is a finite non-Abelian simple group, then  $\rho(G) \leq 3$ . If  $G$  is a group of Lie type over a field  $K$  with  $\text{char } K \neq 2$  and  $K \neq \mathbb{F}_3$ , or a sporadic group not isomorphic to  $Fi_{22}$ ,  $Fi_{23}$ , then  $\rho(G) = 2$ .*

The characterizations of the solvable radical described above should be compared with earlier results. Namely, let  $F_2 = F(x, y)$  denote the free two generator group. Define a sequence  $\vec{e} = e_1, e_2, e_3, \dots$ , where  $e_i(x, y) \in F_2$ , by

$$e_1(x, y) = [x, y], \dots, e_n(x, y) = [e_{n-1}(x, y), y], \dots$$

An element  $g \in G$  is called an Engel element if for every  $a \in G$  there exists  $n = n(a, g)$  such that  $e_n(a, g) = 1$ .

In 1957 R. Baer proved the following theorem [2,13]:

**Theorem 1.11.** *The nilpotent radical of a Noetherian group  $G$  coincides with the collection of all Engel elements of  $G$ .*

In particular, Baer's theorem holds for finite groups. Similar theorems have been established for many classes of infinite groups satisfying some additional conditions (see, for example, [18,17]).

A tempting but difficult problem is to find a counterpart of Baer's theorem for the solvable radical of a finite group. In other words, one has to find an Engel-like sequence  $\vec{u} = \vec{u}_n(x, y)$  such that  $g \in G$  belongs to the solvable radical  $R(G)$  if and only if for every  $a \in G$  there exists  $n = n(a, g)$  such that  $u_n(a, g) = 1$ . First results towards a solution of this problem have been obtained in [4–6,3].

In [12] a Thompson-like characterization of the solvable radical of a finite group has been obtained (it also holds for any linear group or a PI-group):

**Theorem 1.12.** [12] *The solvable radical  $R(G)$  of a finite group  $G$  coincides with the set of all elements  $g \in G$  such that for any  $a \in G$  the subgroup generated by  $g$  and  $a$  is solvable.*

This theorem can be viewed as an implicit description of the solvable radical since it does not provide any explicit formulas which determine if a particular element belongs to  $R(G)$ . The results of the present paper can be viewed as a next step to a more explicit characterization of the solvable radical.

## 2. Theorem 1.8 implies Theorem 1.1

Suppose Theorem 1.8 holds. Let us prove Theorem 1.1.

For brevity, let us call the elements  $g \in G$  satisfying the condition of Theorem 1.1, *suitable*. Clearly, all the elements of the radical  $R(G)$  are suitable. We want to show that any suitable element  $g$  belongs to  $R(G)$ . We may assume that  $G$  is semisimple (i.e.  $R(G) = 1$ ).

Recall that any finite semisimple group  $G$  contains a unique maximal normal centreless completely reducible (CR) subgroup (by definition, CR means a direct product of finite non-Abelian simple groups) which is called the CR-radical of  $G$  (see [19, 3.3.16]). We call a product of the isomorphic factors in the decomposition of the CR-radical an *isotypic component* of  $G$ . Denote the CR-radical of  $G$  by  $V(G)$ . This is a characteristic subgroup of  $G$ .

Let  $G$  be a minimal counterexample to the statement of Theorem 1.1. Then it is easy to show that  $V(G)$  has only one isotypic component. Any  $g \in G$  acts as an automorphism  $\bar{g}$  on  $V(G) = H_1 \times \cdots \times H_n$ , where all  $H_i$ ,  $1 \leq i \leq n$ , are isomorphic simple non-Abelian groups.

Suppose that  $g$  is a suitable element. Using the 'one and a half' generation theorem for simple groups by Guralnick and Kantor [11, Corollary of Th. I on p. 745], one can show that  $\bar{g}$  cannot act on  $V(G)$  as a nontrivial element of the symmetric group  $S_n$ . So we can assume that  $g$  acts as an automorphism of a simple group  $H$ . Consider the extension  $G_1$  of the group  $H$  with the automorphism  $\bar{g}$ . Since  $G_1$  has no centre, one can find  $x \in H$  such that  $[x, \bar{g}] \neq 1$ . By Theorem 1.8 there exist  $y_1, y_2, y_3 \in H$  such that the subgroup  $\langle [[x, g], y_1], [[x, g], y_2], [[x, g], y_3] \rangle$  is not solvable. Using the formula  $y[x, g]y^{-1} = [x, g][[g, x], y]$ , we obtain

$$\langle [[x, g], y_1], [[x, g], y_2], [[x, g], y_3] \rangle \leq \langle g, x^{-1}gx, y_i^{-1}gy_i, y_i^{-1}x^{-1}gx y_i \mid i = 1, 2, 3 \rangle.$$

As  $g$  is suitable, the latter subgroup must be solvable. Contradiction with the choice of  $y_i$ .

**Remark 2.1.** The above argument shows that Conjectures 1.3 and 1.4 follow from the analogue of Theorem 1.8 for the almost simple groups.

## 3. Scheme of the proof of Theorem 1.8

### 3.1. Alternating groups

**Proposition 3.1.** *Let  $G = A_n$ ,  $n \geq 5$ . Then  $\rho(G) = 2$ .*

**Proof.** We proceed by induction. For  $n = 5, 6$  the statement can be checked in a straightforward manner. To make the induction step, we consider separately the cases where  $y$  can (or cannot) be represented in the form  $y = \sigma \tau$ ,  $\sigma \in A_m$ ,  $\sigma \neq 1$ ,  $m < n$ .  $\square$

### 3.2. Groups of Lie type, $\text{char}(K) \neq 2$ and $|K| \neq 3$

Let first  $G$  be a group of rank 1.

**Proposition 3.2.** *Let  $G$  be one of the groups*

$$A_1(q) \ (q \neq 2, 3), \quad {}^2A_2(q^2) \ (q \neq 2), \quad {}^2B_2(2^{2m+1}) \ (m \geq 1), \quad {}^2G_2(3^{2m+1}) \ (m \geq 0).$$

Then  $\rho(G) = 2$ .

The proof involves calculations based on different canonical decompositions in  $G$ . The uniform part of the proof relies on the following theorem of Gow (compare with [7]) regarding conjugacy classes of semisimple elements in Chevalley groups:

**Theorem 3.3.** [10] *Let  $G$  be a finite simple group of Lie type, and let  $g \neq 1$  be a semisimple element in  $G$ . Let  $L$  be a conjugacy class of  $G$  consisting of regular semisimple elements. Then there exist a regular semisimple  $x \in L$  and  $z \in G$  such that  $g = [x, z]$ .*

Certain steps require, however, explicit matrix representations for the groups of rank 1 (see [14,15]). As usual, groups over small fields are considered separately.

Suppose now that the rank of  $G$  is greater than 1.

**Theorem 3.4.** *Let  $G$  be a Chevalley group of rank  $> 1$  over a field  $K$  with  $\text{char } K \neq 2$ ,  $K \neq \mathbb{F}_3$ . Then  $\rho(G) = 2$ .*

We use the Levi decomposition of  $G$  together with arguments from [9] (based on the notions of generalized Bruhat cells and generalized Coxeter elements) in order to reduce to rank 1 case.

### 3.3. Groups of Lie type, $\text{char}(K) = 2$ or $|K| = 3$

**Proposition 3.5.** *Let  $G$  be a non-solvable Chevalley group over a field  $K$ , where either  $\text{char } K = 2$  or  $K = \mathbb{F}_3$ . Then  $\rho(G) \leq 3$ .*

The proof goes by reduction to the case of groups of rank at most 3 and involves more technicalities. In particular, the case  $G = {}^2F_4(q)$  requires separate consideration. Some groups of small ranks are treated by MAGMA.

Note that the estimate of Proposition 3.5 is sharp as follows from the case of groups generated by 3-transpositions (see [8,1] for definitions and notations).

**Proposition 3.6.** *Let  $G$  be one of the following groups (with the notation of [1]): a symmetric group  $S_n$ ; a symplectic group  $\text{Sp}(2n, 2)$  ( $n \geq 2$ ); an orthogonal group  $\text{O}^\mu(2n, 2)$  for  $\mu \in \{-1, 1\}$  and  $n \geq 2$ ; a unitary group  $\text{PSU}(n, 2)$  ( $n \geq 4$ ); an orthogonal group  $\text{O}^{\mu, \pi}(n, 3)$  for  $\mu \in \{-1, 1\}$ ,  $\pi \in \{-1, 1\}$ , and  $n \geq 4$ ; one of Fischer's groups  $Fi_{22}$ ,  $Fi_{23}$ ,  $Fi_{24}$ . Then  $G$  contains a nontrivial 2-radical element.*

### 3.4. Sporadic groups

**Proposition 3.7.** *Let  $G$  be a sporadic simple group. Then  $\rho(G) = 3$  for  $G = Fi_{22}, Fi_{23}$  and  $\rho(G) = 2$  for all the remaining groups.*

For all groups except the Monster  $M$ , our main tool was MAGMA. We relied on the ATLAS classification of conjugacy classes of maximal cyclic subgroups [21]. For larger sporadic groups we had to replace most standard MAGMA procedures by our own ones in order to avoid storing the whole group and large subgroups. In particular, to check whether a subgroup under consideration is not solvable, we used the Hall–Thompson criterion [20]: a group  $H$  is non-solvable if and only if it contains elements  $a, b, c$  of pairwise coprime orders such that  $abc = 1$ .

Our proof for  $M$  also relies on the classification of conjugacy classes. However, all other arguments are theoretic. We reduce the statement to checking the elements  $g$  of prime orders  $p$  each of those (except for  $p = 41$ ) can be included in some proper simple subgroup of  $M$  [16]. The case  $p = 41$  is treated separately using the fact that the normalizer  $N_M(\langle g \rangle) = 41 \cdot 40$  is a maximal subgroup of  $M$  [21].

## Acknowledgement

Kunyavskii and Plotkin were partially supported by the Ministry of Absorption (Israel), the Israeli Science Foundation founded by the Israeli Academy of Sciences – Center of Excellence Program, and the Minerva Foundation through the Emmy Noether Research Institute of Mathematics, and by the RTN network HPRN-CT-2002-00287. A substantial part of this work was done during Gordeev's visits to Bar-Ilan University in May 2005 and May 2006 and the visit of Grunewald, Kunyavskii and Plotkin to MPIM (Bonn) during the activity on "Geometry and Group Theory" in July 2006. The support of these institutions is highly appreciated.

We are very grateful to ~~B.I. Plotkin, J.N. Bray~~ and N.A. Vavilov for useful discussions and correspondence.

## References

- [1] M. Aschbacher, 3-Transposition Groups, Cambridge Univ. Press, Cambridge, 1997.
- [2] R. Baer, Engelsche Elemente Noetherscher Gruppen, Math. Ann. 133 (1957) 256–270.
- [3] T. Bandman, M. Borovoi, F. Grunewald, B. Kunyavskii, E. Plotkin, Engel-like characterization of radicals in finite dimensional Lie algebras and finite groups, Manuscripta Math. 119 (2006) 365–381.
- [4] T. Bandman, G.-M. Greuel, F. Grunewald, B. Kunyavskii, G. Pfister, E. Plotkin, Two-variable identities for finite solvable groups, C. R. Acad. Sci. Paris, Ser. I 337 (2003) 581–586.
- [5] T. Bandman, G.-M. Greuel, F. Grunewald, B. Kunyavskii, G. Pfister, E. Plotkin, Identities for finite solvable groups and equations in finite simple groups, Compositio Math. 142 (2006) 734–764.
- [6] J.N. Bray, J.S. Wilson, R.A. Wilson, A characterization of finite soluble groups by laws in two variables, Bull. London Math. Soc. 37 (2005) 179–186.
- [7] E.W. Ellers, N. Gordeev, On the conjectures of J. Thompson and O. Ore, Trans. Amer. Math. Soc. 350 (1998) 3657–3671.
- [8] B. Fischer, Finite groups generated by 3-transpositions, I, Invent. Math. 12 (1971) 232–246.
- [9] N. Gordeev, J. Saxl, Products of conjugacy classes in Chevalley groups, I: Extended covering numbers, Israel J. Math. 130 (2002) 207–248.
- [10] R. Gow, Commutators in finite simple groups of Lie type, Bull. London Math. Soc. 32 (2000) 311–315.
- [11] R.M. Guralnick, W.M. Kantor, The probability of generating a simple group, J. Algebra 234 (2000) 743–792.
- [12] R. Guralnick, B. Kunyavskii, E. Plotkin, A. Shalev, Thompson-like characterization of radicals in groups and Lie algebras, J. Algebra 300 (2006) 363–375.
- [13] B. Huppert, Endliche Gruppen, I, Springer-Verlag, Berlin, 1979.
- [14] B. Huppert, N. Blackburn, Finite Groups, III, Springer-Verlag, Berlin, 1982.
- [15] G. Kemper, F. Lübeck, K. Magaard, Matrix generators for the Ree groups  ${}^2G_2(q)$ , Comm. Algebra 29 (2001) 407–413.
- [16] S.P. Norton, Anatomy of the Monster, I, in: R. Curtis, R. Wilson (Eds.), The Atlas of Finite Groups: Ten Years on, in: London Math. Soc. Lecture Note Ser., vol. 249, Cambridge Univ. Press, Cambridge, 1998, pp. 198–214.
- [17] V.P. Platonov, Engel elements and radical in PI-algebras and topological groups, Dokl. Akad. Nauk SSSR 161 (1965) 288–291 (in Russian).
- [18] B.I. Plotkin, Notes on Engel groups and Engel elements in groups. Some generalizations, Izv. Ural. Univ. Ser. Mat. Mekh. 36 (7) (2005) 153–166; available at <http://arXiv.org/math.GR/0406100>.
- [19] D.J.S. Robinson, A Course in the Theory of Groups, Springer-Verlag, New York, 1995.
- [20] J. Thompson, Non-solvable finite groups all of whose local subgroups are solvable, Bull. Amer. Math. Soc. 74 (1968) 383–437.
- [21] R.A. Wilson, et al., A world-wide-web atlas of group representations, <http://brauer.maths.qmul.ac.uk/Atlas/>.