

# Word Maps and Word Maps with Constants of Simple Algebraic Groups<sup>1</sup>

N. L. Gordeev<sup>a,b,\*</sup>, B. È. Kunyavskii<sup>c, \*\*</sup>, and E. B. Plotkin<sup>c</sup>

Received June 20, 2016

**Abstract**—In the present paper, we consider word maps  $w: G^m \rightarrow G$  and word maps with constants  $w_\Sigma: G^m \rightarrow G$  of a simple algebraic group  $G$ , where  $w$  is a nontrivial word in the free group  $F_m$  of rank  $m$ ,  $w_\Sigma = w_1\sigma_1w_2 \cdots w_r\sigma_rw_{r+1}$ ,  $w_1, \dots, w_{r+1} \in F_m$ ,  $w_2, \dots, w_r \neq 1$ ,  $\Sigma = \{(\sigma_1, \dots, \sigma_r) \mid \sigma_i \in G \setminus Z(G)\}$ . We present results on the images of such maps, in particular, we prove a theorem on the dominance of “general” word maps with constants, which can be viewed as an analogue of a well-known theorem of Borel on the dominance of genuine word maps. Besides, we establish a relationship between the existence of unipotents in the image of a word map and the structure of the representation variety  $R(\Gamma_w, G)$  of the group  $\Gamma_w = F_m/\langle w \rangle$ .

**DOI:** 10.1134/S1064562416060077

## 1. WORD MAPS WITH CONSTANTS

For any group  $G$  and any non-empty word  $w$  in the free group  $F_m$  of rank  $m$  one can define the *word map*  $w: G^m \rightarrow G$  by the formula  $w((g_1, \dots, g_m)) = w(g_1, \dots, g_m)$  (we substitute the elements  $g_i$  instead of the variables  $x_i$ ). Recently one could observe growing interest to the study of word maps of simple algebraic groups (see [1–3, 5, 8, 9]). For such groups we also consider here word maps with constants. Namely, let  $G$  be a simple algebraic group defined over an algebraically closed field  $K$  (we identify the group  $G$  with the group of points  $G(K)$ ), and let  $w_1, \dots, w_{r+1} \in F_m$ , where  $w_2, \dots, w_r \neq 1$ ,  $\Sigma = \{\sigma_1, \dots, \sigma_r \mid \sigma_i \in G \setminus Z(G)\}$  (we allow  $\sigma_i = \sigma_j$  for  $i \neq j$ ). The expression  $w_\Sigma = w_1\sigma_1w_2 \cdots w_r\sigma_rw_{r+1}$  is called a word with constants (we regard usual words as words with constants by setting  $\Sigma = \emptyset$ ,  $w = w_1$ ). The behaviour of words with constants on simple algebraic groups was studied, in particular, in [6, 7, 15]. A word with constants also gives rise to a natural word map with constants  $w_\Sigma: G^m \rightarrow G$ . In [6] such maps were used for studying products of conjugacy classes, and in [8] they served as a method for studying genuine word maps.

One of the main questions of the theory of word maps concerns their surjectivity (the answer is unknown even for the group  $G = \mathrm{SL}_2(C)$ , see [9]). According to a theorem of A. Borel [4], the word maps

<sup>1</sup> The article was translated by the authors.

<sup>a</sup>Herzen State Pedagogical University of Russia, St. Petersburg

<sup>b</sup>St. Petersburg State University

<sup>c</sup>Bar-Ilan University, Ramat Gan, Israel

\*e-mail: nickgordeev@mail.ru

\*\*e-mail: kunyav@gmail.com

of the simple algebraic groups are dominant, i.e., the image  $\mathrm{Im} w$  of such a map contains a dense open subset of  $G$ . A word map with constants is not necessarily dominant (for example, for  $w_\Sigma = x\sigma x^{-1}$ ). However, for a “general” word with constants such a map turns out to be dominant.

**Theorem 1.** *Let  $\Omega_r = (w_1, \dots, w_{r+1})$  be a sequence of words from  $F_m$  where  $w_2, \dots, w_r \neq 1$ . Suppose that  $\prod_{i=1}^{r+1} w_i \notin [F_m, F_m]$ . Then there is a non-empty Zariski open subset  $U(\Omega_r) \subset G^r$  such that for every sequence  $\Sigma = (\sigma_1, \dots, \sigma_r) \in U(\Omega_r)$  the map  $w_\Sigma: G^m \rightarrow G$  is dominant.*

We also consider the case of word maps with constants for which we have  $\prod_{i=1}^{r+1} w_i = 1$ . Namely, let  $w(x, y) \in F_2$  and  $\sigma \in G$ . Then the map  $w_\sigma: G \rightarrow G$  defined by the formula  $w_\sigma(x) = w(x, \sigma)$  is a word map with constants (here the constants are powers of  $\sigma$ ). We have the following theorem, which can serve as a tool in studying word maps in two variables (see [8]).

**Theorem 2.** *Let  $w \in [F_2, F_2]$ . Then there is a non-empty Zariski open subset  $U(w) \subset G$  such that for every  $\sigma \in U(w)$  the set  $\{g(\mathrm{Im} w_\sigma)g^{-1} \mid g \in G\}$  is Zariski dense in  $G$ .*

## 2. SEMISIMPLE ELEMENTS IN $\mathrm{Im} w$

In the paper [3] it was proven that for  $G = \mathrm{SL}_2(K)$  the image of the word map  $w: \mathrm{SL}_2(K)^m \rightarrow \mathrm{SL}_2(K)$  contains all semisimple elements of  $\mathrm{SL}_2(K)$  (here  $K$  is an algebraically closed field) except possibly  $-\mathbf{1}$  ( $\mathbf{1}$  denotes the identity matrix). Using the fact that for all simple algebraic groups except those of types  $A_r$ ,  $D_{2r+1}$ ,  $E_6$ , the corresponding root system contains a subsystem of the same rank which consists of the union of

disjoint subsystems of type  $A_1$ , one can get the following assertion.

**Theorem 3.** *Let  $G$  be a simple algebraic group, and let  $w: G^m \rightarrow G$  be a word map. Suppose that  $G$  is not of type  $A_r$ ,  $r > 1$ ,  $D_{2r+1}$ , or  $E_6$ . Then every regular semisimple element of  $G$  is contained in  $\text{Im } w$ . Moreover, for every semisimple element  $g \in G$  there exists an element  $g_0$  of order two such that  $gg_0 \in \text{Im } w$ .*

### 3. UNIPOTENT ELEMENTS IN $\text{Im } w$ AND THE REPRESENTATION VARIETY OF A FINITELY GENERATED GROUP

Let  $T$  and  $W$  denote, respectively, a fixed maximal torus and the Weyl group of  $G$ , and let  $\pi: G \rightarrow T/W$  be the quotient morphism (see [14]). For a word map  $w: G^m \rightarrow G$  define  $Y_w = w^{-1}(\mathbf{1})$ ,  $\Xi_w = (\pi \cdot w)^{-1}(\mathbf{1})$  (here  $\mathbf{1}$  denotes the identity element of  $G$  and also the image of the identity element of  $T$  in  $T/W$ ). Then  $Y_w \subset \Xi_w$  are affine subvarieties of  $G^m$ ,  $\Xi_w = \{(g_1, \dots, g_m) \in G^m \mid w(g_1, \dots, g_m) \text{ is a unipotent element}\}$ ,  $Y_w = R(\Gamma_w, G)$  is the variety of representations of the one-relator group  $\Gamma_w = F_m/\langle w \rangle$  in the group  $G$ . Thus, the existence of nontrivial unipotent elements in  $\text{Im } w$  is equivalent to the inequality  $Y_w \neq \Xi_w$ . For example, in the simplest case  $G = \text{SL}_2(K)$  and  $m = 2$  all irreducible components of the variety  $\Xi_w$  are of dimension 5, and thus the existence of nontrivial unipotent elements in  $\text{Im } w$  follows from the existence of irreducible components of  $Y_w$  of dimension  $\leq 4$ .

The representation variety of a group is an important object which can be regarded from various points of view (see, e.g., [10–13]) and may be crucial for answering the question on the existence of unipotent elements in  $\text{Im } w$ .

The existence of unipotent elements in  $\text{Im } w$  is an open question even in the case  $G = \text{SL}_2(K)$ . In [3], Bandman and Zarhin proved that for  $G = \text{SL}_2(K)$  ( $\text{ch } K = 0$ ) and  $w \notin [[F_m, F_m], [F_m, F_m]]$ , the set  $\text{Im } w$  contains all unipotent elements. Besides, they gave an example of a computer-aided calculation for a word  $w \in [[F_m, F_m], [F_m, F_m]]$  such that  $\text{Im } w$  also contains all unipotent elements. We consider a similar example for which we calculate (without using computer) the varieties  $Y_w, \Xi_w$ . Let  $K$  be an algebraically closed field ( $\text{ch } K = 0$ ), and let  $w: \text{SL}_2(K)^2 \rightarrow \text{SL}_2(K)$  be the word map induced by the word  $w(x, y) = [[x, y], x[x, y]x^{-1}]$ . Let  $B$  and  $T$  denote, respectively, the upper triangular and diagonal matrices in  $\text{SL}_2(K)$ , let  $\omega = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ , and let  $C_\omega$  be the conjugacy class of  $\omega$  in  $\text{SL}_2(K)$ . Then we have the following fact.

**Theorem 4.** *The variety  $\Xi_w$  has exactly three irreducible components:*

$$\Xi_w^0 = \overline{\{g(B \times B)g^{-1} \mid g \in G\}},$$

$$\Xi_w^1 = \overline{\{g(T \times \omega B)g^{-1} \mid g \in G\}},$$

$$\Xi_w^2 = C_\omega \times G,$$

and the variety  $Y_w$  also has exactly three irreducible components:

$$Y_w^0 = \Xi_w^0, Y_w^1 = \overline{\{g(T \times \omega T)g^{-1} \mid g \in G\}} \subset \Xi_w^1,$$

$$\dim Y_w^1 = 4, \quad Y_w^2 = \Xi_w^2$$

(here bar stands for the Zariski closure).

Thus the existence of the component  $Y_w^1$  of dimension 4 guarantees that all unipotents lie in  $\text{Im } w$ . Although a full proof of this theorem requires significant technical arguments, the mere fact that all unipotents belong to  $\text{Im } w$  is proved by an elementary calculation of the value of  $w(s, \omega b)$  for  $s \in T$ ,  $s^4 \neq \mathbf{1}$ ,  $b \in B$ ,  $b \notin T$ .

The research of the first author was supported by the Ministry of Education and Science of the Russian Federation and RFBR grant 14–01–00820. The research of the second author was supported by ISF grant 1207/12. The research of the third author was supported by ISF grant 1207/12 and the Emmy Noether Institute for Mathematics.

### REFERENCES

1. T. Bandman and Sh. Garion, *Cent. Eur. J. Math.* **12** (2), 175–211 (2014).
2. T. Bandman and B. Kunyavskii, *J. Algebra* **382**, 282–302 (2013).
3. T. Bandman and Yu. Zarhin, ArXive 1407.3447v4[math.GR]; *Eur. J. Math.* (2016) (doi 10.1007/s40879-016-0101-9); <http://link.springer.com/article/10.1007/s40879-016-0101-9>
4. A. Borel, *Enseign. Math.* **29** (1–2), 151–164 (1983).
5. A. Elkasapy and A. Thom, *Indiana Univ. Math. J.* **63** (5), 1553–1565 (2014).
6. N. Gordeev, *J. Algebra* **173**, 715–744 (1995).
7. N. Gordeev, *St. Petersburg Math. J.* **9** (4), 709–723 (1998).
8. N. Gordeev, *J. Algebra* **425**, 215–244 (2015).
9. E. Klimenko, B. Kunyavskii, J. Morita, and E. Plotkin, *Toyama Math. J.* **37**, 25–53 (2015).
10. A. Lubotzky and A. R. Magid, *Mem. Am. Math. Soc.* **58** (336), 117 (1985).
11. V. P. Platonov, “Rings and manifolds of representations of finitely generated groups,” in *Issues of Algebra* (Universitetskoe, Minsk, 1989), Vol. 4, pp. 36–40 [in Russian].
12. V. P. Platonov and V. V. Benyash-Krivetz, *Proc. Steklov Inst. Math.* **183** (4), 203–213 (1991).
13. A. S. Rapinchuk, V. V. Benyash-Krivetz, and V. I. Chemousov, *Israel J. Math.* **93**, 29–71 (1996).
14. T. A. Springer and R. Steinberg, *Lect. Notes Math.* **131** (1970).
15. A. Stepanov, *J. Algebra* **362**, 12–29 (2012).